



B. TECH DEGREE EXAMINATIONS: NOV/DEC 2022

(Regulation 2018)

Fifth Semester

FASHION TECHNOLOGY

U18MAT5102: Probability and Statistics

(Statistical Tables required)

COURSE OUTCOMES

CO1:	Compute measures of central tendencies, dispersion, and correlation between variables, and predict unknown values using regression.
CO2:	Understand and apply the concept of probability and random variables.
CO3:	Construct probabilistic models for observed phenomena through distributions, which play an important role in many engineering applications.
CO4:	Perform hypothesis testing and interpret the results.
CO5:	Understand the principles of design of experiments and perform analysis of variance.
CO6:	Sketch control charts and comment on the process control.

Time: Three Hours

Maximum Marks: 100

Answer all the Questions: -

PART A (10 x 2 = 20 Marks)
(Answer not more than 40 words)

1.	If the marks of 5 students are 45, 92, 26, 81 and 72, find the range and coefficient of range.	CO1	[K ₂]																				
2.	Calculate the rank correlation coefficient between marks in the selection test (X) and the proficiency test (Y) of 9 recruits. <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td>X:</td> <td>10</td> <td>15</td> <td>12</td> <td>17</td> <td>13</td> <td>16</td> <td>24</td> <td>14</td> <td>22</td> </tr> <tr> <td>Y:</td> <td>30</td> <td>42</td> <td>45</td> <td>46</td> <td>33</td> <td>34</td> <td>40</td> <td>35</td> <td>39</td> </tr> </table>	X:	10	15	12	17	13	16	24	14	22	Y:	30	42	45	46	33	34	40	35	39	CO1	[K ₃]
X:	10	15	12	17	13	16	24	14	22														
Y:	30	42	45	46	33	34	40	35	39														
3.	Four persons are chosen at random from a group containing 3 men, 2 women and 4 children. Find the probability that exactly two of them will be children.	CO2	[K ₂]																				
4.	Find the moment generating function of the distribution whose probability mass function is $p(x) = \begin{cases} q^x p, & x = 0, 1, 2, \dots \\ 0, & \text{otherwise} \end{cases}$	CO2	[K ₄]																				
5.	For usual notation with binomial variate X, find p when n = 6 and $9P(x = 4) = P(x = 2).$	CO3	[K ₃]																				

6.	A manufacturer of cotter pins knows that 5% of his product is defective. If he sells pins in boxes of 100 and guarantees that not more than 10 pins will be defective, what is the approximate probability that a box will fail to meet the guaranteed quality.	CO3	[K ₃]																								
7.	A simple sample of heights of 6400 Englishmen has a mean of 170 cm and a S.D. of 6.4 cm, while a simple sample of heights of 1600 Americans has a mean of 172 cm and a S.D. of 6.3 cm. Find the test statistic.	CO4	[K ₂]																								
8.	A sample of 200 persons with a particular disease was selected. Out of these, 100 were given a drug and the others were not given any drug. The results are as follows: <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>No. of persons</td> <td>Drug</td> <td>No drug</td> </tr> <tr> <td>Cured</td> <td>65</td> <td>55</td> </tr> <tr> <td>Not cured</td> <td>35</td> <td>45</td> </tr> </table> <p>Find the test statistic.</p>	No. of persons	Drug	No drug	Cured	65	55	Not cured	35	45	CO4	[K ₂]															
No. of persons	Drug	No drug																									
Cured	65	55																									
Not cured	35	45																									
9.	A random sample is selected from each of 3 makes of ropes and their breaking strength are measured with the following results. <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>I</td> <td>70</td> <td>72</td> <td>75</td> <td>80</td> <td>83</td> <td></td> <td></td> </tr> <tr> <td>II</td> <td>100</td> <td>110</td> <td>108</td> <td>112</td> <td>113</td> <td>120</td> <td>107</td> </tr> <tr> <td>III</td> <td>60</td> <td>65</td> <td>57</td> <td>84</td> <td>87</td> <td>73</td> <td></td> </tr> </table> <p>Find total sum of squares.</p>	I	70	72	75	80	83			II	100	110	108	112	113	120	107	III	60	65	57	84	87	73		CO5	[K ₂]
I	70	72	75	80	83																						
II	100	110	108	112	113	120	107																				
III	60	65	57	84	87	73																					
10.	The following data represents number of end breaks (per 100 spindle hours) observed on a ring frame on eight different days. <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>Day no.</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> <td>7</td> <td>8</td> </tr> <tr> <td>Number of end breaks</td> <td>2</td> <td>8</td> <td>3</td> <td>2</td> <td>4</td> <td>6</td> <td>5</td> <td>7</td> </tr> </table> <p>Find Control limit (CL) for c-chart.</p>	Day no.	1	2	3	4	5	6	7	8	Number of end breaks	2	8	3	2	4	6	5	7	CO6	[K ₃]						
Day no.	1	2	3	4	5	6	7	8																			
Number of end breaks	2	8	3	2	4	6	5	7																			

**Answer any FIVE Questions: -
PART B (5 x 16 = 80 Marks)
(Answer not more than 400 words)**

11.	a)	Marks obtained by 10 students in Mathematics (X) and Statistics (Y) are given below: <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>X:</td> <td>60</td> <td>34</td> <td>40</td> <td>50</td> <td>45</td> <td>40</td> <td>22</td> <td>43</td> <td>42</td> </tr> <tr> <td>Y:</td> <td>75</td> <td>32</td> <td>33</td> <td>40</td> <td>45</td> <td>33</td> <td>12</td> <td>30</td> <td>34</td> </tr> </table> <p>Find the two regression lines. Also find y when x = 55.</p>	X:	60	34	40	50	45	40	22	43	42	Y:	75	32	33	40	45	33	12	30	34	8	CO1	[K ₃]
X:	60	34	40	50	45	40	22	43	42																
Y:	75	32	33	40	45	33	12	30	34																

	b)	Calculate mean, median and mode for the following data:	8	CO1	[K ₄]																
		<table border="1"> <tr> <td>Interval:</td> <td>0-10</td> <td>10-20</td> <td>20-30</td> <td>30-40</td> <td>40-50</td> <td>50-60</td> <td>60-70</td> </tr> <tr> <td>Frequency:</td> <td>7</td> <td>7</td> <td>14</td> <td>17</td> <td>15</td> <td>8</td> <td>2</td> </tr> </table>	Interval:	0-10	10-20	20-30	30-40	40-50	50-60	60-70	Frequency:	7	7	14	17	15	8	2			
Interval:	0-10	10-20	20-30	30-40	40-50	50-60	60-70														
Frequency:	7	7	14	17	15	8	2														
12.	a)	In a bolt factory, machines A, B and C manufacture respectively 25%, 35% and 40% of the total. Of their output 5, 4, 2 percent are defective bolts. A bolt is drawn at random from the product and is found to be defective. What are the probabilities that it was manufactured by machines A, B and C?	8	CO2	[K ₄]																
	b)	Find the mean, variance of the random variable X which has the following density function $f(x) = \begin{cases} x, & 0 < x < 1 \\ 2 - x, & 1 < x < 2 \\ 0, & \text{otherwise.} \end{cases}$	8	CO2	[K ₄]																
13.	a)	In a test of 2000 electric bulbs, it was found that the life of a particular make was normally distributed with an average life of 2040hrs and SD of 60 hrs. Estimate (i) The number of bulbs likely to burn for (i) more than 2150hrs(ii) less than 1950hrs and (iii) more than 1920 hrs but less than 2160hrs.	8	CO3	[K ₃]																
	b)	Fit a binomial distribution for the following data <table border="1"> <tr> <td>x:</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> </tr> <tr> <td>f:</td> <td>5</td> <td>18</td> <td>28</td> <td>12</td> <td>7</td> <td>6</td> <td>4</td> </tr> </table>	x:	0	1	2	3	4	5	6	f:	5	18	28	12	7	6	4	8	CO3	[K ₄]
x:	0	1	2	3	4	5	6														
f:	5	18	28	12	7	6	4														
14.	a)	Two random samples gave the following results: <table border="1"> <tr> <th>Sample</th> <th>Size</th> <th>Sample mean</th> <th>Sum of the square of deviations from the mean</th> </tr> <tr> <td>1</td> <td>10</td> <td>15</td> <td>90</td> </tr> <tr> <td>2</td> <td>12</td> <td>14</td> <td>108</td> </tr> </table> <p>Examine whether the samples come from the same normal population at 5% level of significance.</p>	Sample	Size	Sample mean	Sum of the square of deviations from the mean	1	10	15	90	2	12	14	108	8	CO4	[K ₄]				
Sample	Size	Sample mean	Sum of the square of deviations from the mean																		
1	10	15	90																		
2	12	14	108																		
	b)	In a sample of 1000 people in Mumbai, 540 are rice eaters and the rest are wheat eaters. Can we assume that both rice and wheat are equally popular in this state at 1% level of significance?	8	CO4	[K ₃]																

15.	a)	<p>The following data represents the number of units of productions per day turned out by different workers using four different types of machines.</p> <table border="1" data-bbox="248 226 1177 495"> <thead> <tr> <th colspan="2"></th> <th colspan="4">Machine Type</th> </tr> <tr> <th colspan="2"></th> <th>A</th> <th>B</th> <th>C</th> <th>D</th> </tr> </thead> <tbody> <tr> <td></td> <td>1</td> <td>44</td> <td>38</td> <td>47</td> <td>36</td> </tr> <tr> <td>Workers</td> <td>2</td> <td>46</td> <td>40</td> <td>52</td> <td>43</td> </tr> <tr> <td></td> <td>3</td> <td>34</td> <td>36</td> <td>44</td> <td>32</td> </tr> <tr> <td></td> <td>4</td> <td>43</td> <td>38</td> <td>46</td> <td>33</td> </tr> <tr> <td></td> <td>5</td> <td>38</td> <td>42</td> <td>49</td> <td>39</td> </tr> </tbody> </table> <p>Test whether the 5 men differ with respect to mean productivity and test whether the mean productivity is the same for the four different machine types.</p>			Machine Type						A	B	C	D		1	44	38	47	36	Workers	2	46	40	52	43		3	34	36	44	32		4	43	38	46	33		5	38	42	49	39	8	CO5	[K ₅]
		Machine Type																																													
		A	B	C	D																																										
	1	44	38	47	36																																										
Workers	2	46	40	52	43																																										
	3	34	36	44	32																																										
	4	43	38	46	33																																										
	5	38	42	49	39																																										
	b)	<p>The following data give the average life in hours and range in hours of 12 samples each of 5 lamps. Construct the control charts for \bar{X} and R and comment on the state of control.</p> <table border="1" data-bbox="248 779 1312 898"> <tbody> <tr> <td>\bar{X}</td> <td>120</td> <td>127</td> <td>152</td> <td>157</td> <td>160</td> <td>134</td> <td>137</td> <td>123</td> <td>140</td> <td>144</td> <td>120</td> <td>127</td> </tr> <tr> <td>R</td> <td>30</td> <td>44</td> <td>60</td> <td>34</td> <td>38</td> <td>35</td> <td>45</td> <td>62</td> <td>39</td> <td>50</td> <td>35</td> <td>41</td> </tr> </tbody> </table>	\bar{X}	120	127	152	157	160	134	137	123	140	144	120	127	R	30	44	60	34	38	35	45	62	39	50	35	41	8	CO6	[K ₄]																
\bar{X}	120	127	152	157	160	134	137	123	140	144	120	127																																			
R	30	44	60	34	38	35	45	62	39	50	35	41																																			
16.	a)	<p>Two researchers adopted different sampling techniques while investigating the same group of students to find the number of students falling into different intelligence level. The results are as follows:</p> <table border="1" data-bbox="248 1140 1149 1419"> <thead> <tr> <th>Research</th> <th>Below average</th> <th>Average</th> <th>Above average</th> <th>Excellent</th> <th>Total</th> </tr> </thead> <tbody> <tr> <td>X</td> <td>86</td> <td>60</td> <td>44</td> <td>10</td> <td>200</td> </tr> <tr> <td>Y</td> <td>40</td> <td>33</td> <td>25</td> <td>2</td> <td>100</td> </tr> <tr> <td>Total</td> <td>126</td> <td>93</td> <td>69</td> <td>12</td> <td>300</td> </tr> </tbody> </table> <p>Would you say that the sampling techniques adopted by the two researchers are significantly different?</p>	Research	Below average	Average	Above average	Excellent	Total	X	86	60	44	10	200	Y	40	33	25	2	100	Total	126	93	69	12	300	8	CO4	[K ₄]																		
Research	Below average	Average	Above average	Excellent	Total																																										
X	86	60	44	10	200																																										
Y	40	33	25	2	100																																										
Total	126	93	69	12	300																																										
	b)	<p>The prices of two commodities over 10 weeks are given below. Find out which price shows less variation.</p> <table border="1" data-bbox="248 1686 1312 1797"> <tbody> <tr> <td>A:</td> <td>54</td> <td>55</td> <td>53</td> <td>56</td> <td>52</td> <td>52</td> <td>58</td> <td>49</td> <td>50</td> <td>51</td> </tr> <tr> <td>B:</td> <td>108</td> <td>107</td> <td>105</td> <td>106</td> <td>105</td> <td>103</td> <td>102</td> <td>104</td> <td>104</td> <td>101</td> </tr> </tbody> </table>	A:	54	55	53	56	52	52	58	49	50	51	B:	108	107	105	106	105	103	102	104	104	101	8	CO1	[K ₃]																				
A:	54	55	53	56	52	52	58	49	50	51																																					
B:	108	107	105	106	105	103	102	104	104	101																																					
